



Combining Structure- and Ligand-Based Approaches to Model Receptor-Mediated Toxic Effects



Markus Lill

Department of Medicinal Chemistry
and Molecular Pharmacology

Purdue University



Overview





Introduction

Methodology

Applications


Current directions






Biological systems

Nuclear receptors




Cytochrome P450 enzymes







Endogenous ligands




17- β estradiol (ER)



testosterone (AR)




progesterone (PR)



cortisol (GR)




triiodothyronine (TR)




1,25-dihydroxy vitamin D₃ (VDR)

- small (volume = 250-350 Å³)
- rigid
- highly hydrophobic
- polar groups predominantly at the ends

selective for very potent ligand molecules (nM):



2.2 nM (AR)




385 nM (AR)


not selective for moderate-binding ligand molecules (μ M):




6 μ M (AR)




11 μ M (AR)




Receptor-mediated toxicity







NR & P450

NR induce CYP genes







Specific factors



Overview





Introduction

Methodology

Applications

Current directions






Prediction of binding affinities

BIG challenge: Predict binding affinities in reasonable amount of time


Docking & Scoring




~1min

Computing time

Simulations (FEP)




1day - 1week




Quantification of binding affinities


Docking




Binding modes










Protein modeling based on QSAR



Quantification



QSAR




Training set

Test set



Alignment




Optimize model




Predict affinities






Quantification of binding affinities


Docking




Binding modes



Protein modeling based on QSAR




Quantification







Issues:


- No unique docking solution



Bioactive conformation



Coumestrol
binding to the
estrogen
receptor






Quantification of binding affinities


Docking




Binding modes



Protein modeling based on QSAR




Quantification




Issues:

- No unique docking solution
- Induced protein fit





Induced fit




HIV-1 RT with
nevirapine


1051U91



AAP



HEPT






Quantification of binding affinities


Docking




Binding modes



QSAR




Quantification





Issues:

- No unique docking solution
- Induced protein fit



Flexible docking



1. Autodock ligand to binding site




2. Monte Carlo search (flexible minimization & dynamic solvation)



3. Identify new binding mode





N best solutions → 4D data set




Quantification of binding affinities


Docking




Binding modes



QSAR





Quantification



Issues:


- No unique docking solution
- Induced protein fit






Quasar


3D-QSAR:
Simulation
based on
3D
structures



5D-QSAR:
Ensemble of
induced-fit
scenarios




4D-QSAR: Representation by an ensemble of conformations




Scoring function based on a directional force field and explicit desolvation:

$$\Delta G = \Delta E_{\text{force field}} - \Delta G_{\text{solvation}} - T\Delta S - \Delta G_{\text{induced fit}}$$




Induced fit

Estrogen receptor




17 β -estradiol

Raloxifene




Induced fit

Neuraminidase




Zanamavir

2-
dihydropyran-
phenethyl-
propyl-
carboxamide
(2-PPC)






Raptor


Dual-shell representation



Adaptation due to ligand properties



$$HB_{\alpha} = (HBA_{\alpha}^0 - HBD_{\alpha}^0) \frac{1}{1 + \exp(\delta_{hb,\alpha}(hb_{\alpha} - \omega_{hb,\alpha}))} + HBD_{\alpha}^0$$




Raptor


Markus Lill

PURDUE
UNIVERSITY

Dual-shell representation



Adaptation due to ligand properties





$$HB_{\alpha} = (HBA_{\alpha}^0 - HBD_{\alpha}^0) \frac{1}{1 + \exp(\delta_{hb,\alpha}(hb_{\alpha} - \omega_{hb,\alpha}))} + HBD_{\alpha}^0$$

Scoring function based on hydrophobicity and hydrogen bonding

$$\Delta G = \Delta G_{\text{const}} + \Delta G_{\text{HPhob}} + \Delta G_{\text{HBond}} - \Delta G_{\text{induced fit}} - T\Delta S$$

Overview





Introduction

Methodology

Applications

Current directions







Applications

Nuclear receptors


- Estrogen receptors
- Androgen receptor
- Thyroid receptors






Estrogen receptor



116 substances
8 substance classes




DES and related compounds




antiestrogens




steroids




α - α -dimethyl-b-ethyl
allenolic acid




phytoestrogens



diphenylmethanes






biphenyls



phenols


Estrogen receptor



$$q^2 = 0.908$$

$$p^2 = 0.907$$


Lill, M.A.; Vedani, A.; Dobler, M. *J. Med. Chem.* **2004**, *47*, 6174-6186.




Applications

Nuclear receptors


- Estrogen receptors
- Androgen receptor
- Thyroid receptors






Androgen receptor

steroids




119 compounds


6 substance classes




DES and related compounds




phytoestrogens





diphenylmethanes




phenols






organochlorines






Androgen receptor




$$q^2 = 0.858$$

$$p^2 = 0.792$$






Androgen receptor

polychlorinated biphenyls




3,4-diphenyltetrahydrofuran







Applications

Cytochrome P450
• CYP 3A4







Cytochrome P450 3A4





acetaminophen




amiodarone




vinblastine




clotrimazole




alprazolam




benzo(a)pyrene



Cytochrome P450 3A4






Cytochrome P450 3A4


amiodarone:

N-deethylation




dapsone:

N-hydroxylation




diltiazem:


N-demethylation



quinine:




3-hydroxylation






Cytochrome P450 3A4

Different binding modes



terfenadine



Cytochrome P450 3A4



Markus Lill

PURDUE
UNIVERSITY

48 compounds
(192 representations)



$q^2 = 0.825$

$p^2 = 0.659$



Lill, M.A.; Dobler, M.; Vedani, A. *ChemMedChem* **2006**, 1, 73-81.

Overview





Introduction

Methodology

Applications

Current directions







Current directions


Current limitations and research directions

- Flexibility during docking limited to side-chain motion





Specific factors





Agonism/Antagonism





Agonism/Antagonism

Androgen receptor

hydroxyflutamide




antagonist

R-3



agonist



Markus Lill

PURDUE
UNIVERSITY

Acknowledgements

Department of Medicinal Chemistry
Purdue University



Matt Danielson


Department of Pharmacy
University of Basel
&
Biographics Laboratory 3R



Angelo Vedani



Beat Ernst



Max Dobler



Fabienne
Winiger

Purdue University Start-Up
Swiss National Science Foundation Fellowship (NFP 50)
Jacques en Dolly Gazan Foundation Fellowship